

AD-A239 173



AD _____

2

ETIOLOGY AND RAPID DIAGNOSIS OF HUMAN VIRAL GASTROENTERITIS

Annual Report

Neil R. Blacklow, M.D.

May 24, 1991

(For the period 1 May 1990 - 30 April 1991)

Supported by

U. S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
Fort Detrick, Frederick, Maryland 21702-5012

Contract No. DAMD17-86-C-6196

University of Massachusetts Medical School
Worcester, Massachusetts 01655

Approved for public release; distribution unlimited

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DTIC
ELECTE
AUG 6 1991
S B D

91-06851



91 8 05 02

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION <u>Unclassified</u>			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			4. PERFORMING ORGANIZATION REPORT NUMBER(S)		
5. MONITORING ORGANIZATION REPORT NUMBER(S)			6a. NAME OF PERFORMING ORGANIZATION University of Massachusetts Medical School		
6b. OFFICE SYMBOL (If applicable)			7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Worcester, Massachusetts 01655			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Medical Research & Development Command			8b. OFFICE SYMBOL (If applicable)		
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, Maryland 21702-5012			9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract No. DAMD17-86-C-6196		
10. SOURCE OF FUNDING NUMBERS			11. TITLE (Include Security Classification) Etiology and Rapid Diagnosis of Human Viral Gastroenteritis		
PROGRAM ELEMENT NO. 62787A			PROJECT NO. 3M1- 62787A870		
TASK NO. AE			WORK UNIT ACCESSION NO. WUDA310384		
12. PERSONAL AUTHOR(S) Neil R. Blacklow, M.D.					
13a. TYPE OF REPORT Annual Report		13b. TIME COVERED FROM <u>5/1/90</u> TO <u>4/30/91</u>		14. DATE OF REPORT (Year, Month, Day) 1991 May 24	
15. PAGE COUNT 19					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Enteric Adenoviruses, Astroviruses, Caliciviruses, Enterics, Norwalk Virus, Hybridomas, Infectious Diseases, Monoclonals, Rapid Identification, Snow Mountain Virus, RAI		
06	03				
06	13				
19. ABSTRACT (Continue on reverse if necessary and identify by block number) This project is designed to assess the etiology and establish the rapid diagnosis of human viral gastroenteritis. During the contract year for this report, we have continued to demonstrate the involvement of astroviruses in gastroenteritis. In a previously reported Thai study, we have now determined that the clinical symptoms of patients with astrovirus infections were similar to the symptoms of those with rotavirus infection. Additional studies in Guatemala (5,234 samples) and Australia (1,343 samples) have confirmed the importance of astroviruses in gastroenteritis in comparison to other virus infections (rotaviruses and enteric adenoviruses). During this contract year we also developed monoclonal antibodies to calicivirus and we are evaluating these antibodies for use as diagnostic reagents. Ongoing collaborative epidemiological studies have continued to be done with the U.S. military in overseas populations, utilizing the immunodiagnostic techniques that we have developed for gastroenteritis viruses.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Mrs. Virginia Miller			22b. TELEPHONE (Include Area Code) 301-663-7325		22c. OFFICE SYMBOL SGRD-RMI-S

SUMMARY

This project is designed to assess the etiology and establish the rapid diagnosis of human viral gastroenteritis. Of primary importance is the development and utilization of immunoassays to detect various viruses, with preparation and use of monoclonal antibody reagents where possible. Following the successful development of diagnostic assays, the medical and epidemiological importance of these viruses is being ascertained. During the contract year for this report, we have continued to demonstrate the involvement of astroviruses in gastroenteritis. In a previously reported Thai study, we have now determined that the clinical symptoms of patients with astrovirus infections were similar to the symptoms of those with rotavirus infection. Additional studies in Guatemala (5,234 samples) and Australia (1,343 samples) have confirmed the importance of astroviruses in gastroenteritis in comparison to other virus infections (rotaviruses and enteric adenoviruses). The large number of samples examined was possible because of the monoclonal antibody immunoassays we have developed to rotaviruses and enteric adenoviruses as well as to astroviruses. During this contract year we also developed monoclonal antibodies to calicivirus and we are evaluating these antibodies for use as diagnostic reagents. Ongoing collaborative epidemiological studies have continued to be done with the U.S. military in overseas populations, utilizing the immunodiagnostic techniques that we have developed for gastroenteritis viruses.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Distribution/	
Availability Codes	
Avail and/or	
Dist	Avail
A-1	

FOREWORD

In conducting the research described in this report, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (DHEW Publication No. (NIH) 78-23, Revised 1978).

For the protection of human subjects the investigator(s) have adhered to policies of applicable Federal Law 45CFR46.

CONTENTS

	<u>PAGE</u>
SUMMARY.....	1
FOREWORD.....	2
BACKGROUND INFORMATION ON VIRAL GASTROENTERITIS.....	4
ASTROVIRUS ETIOLOGY OF VIRAL GASTROENTERITIS.....	6
CALICIVIRUS MONOCLONAL ANTIBODIES.....	9
SNOW MOUNTAIN VIRUS MONOCLONAL ANTIBODIES.....	9
ANTIIDIOTYPIC ANTIBODIES AS SURROGATE ANTIGENS.....	9
TYPE-SPECIFIC ASTROVIRUS IDENTIFICATION.....	10
COLLABORATIVE EPIDEMIOLOGICAL STUDIES WITH THE MILITARY.....	10
LITERATURE CITED.....	12
PUBLICATIONS OF WORK SUPPORTED BY THIS CONTRACT.....	15
DISTRIBUTION LIST.....	19

BACKGROUND INFORMATION ON VIRAL GASTROENTERITIS

Acute viral gastroenteritis is an extremely common illness that affects all age groups and occurs in both epidemic and endemic forms (1). It is second in frequency only to the common cold among illnesses affecting United States families under epidemiological surveillance. It is also responsible for some of the common travelers' diarrhea encountered in Latin America, Africa, and Asia. The illness varies in its clinical presentation, but in general it begins with an explosive onset, and consists of varying disabling combinations of diarrhea, nausea, vomiting, low grade fever, abdominal cramps, headache, anorexia, myalgia, and malaise. It can be severe, indeed fatal, in the elderly, infant, debilitated, or malnourished patient.

Viral gastroenteritis occurs primarily in two epidemiologically distinct clinical forms (1). One entity is characteristically epidemic and is responsible for family and community-wide outbreaks of gastroenteritis among older children and adults. In recent years, one agent, Norwalk virus, has been shown to be responsible for about 40 percent of these disease outbreaks in the United States. Other Norwalk-like viruses have also been discovered such as Hawaii virus and Snow Mountain virus, and although they have not been well studied epidemiologically, they are likely to be responsible for many more epidemic cases of this illness.

The second clinical entity is usually sporadic, occasionally epidemic, and occurs predominantly in infants and young children (1). However, as noted below it can occur in adults. This form of illness typically produces severe diarrhea that commonly lasts for five to eight days and is usually accompanied by fever and vomiting. Rotavirus is responsible for nearly one half of the cases of this clinical entity requiring hospitalization. Although the major target of rotavirus is the very young, it can produce surprisingly severe clinical disease in adults (1,2).

Breakthroughs in determining the medical importance of Norwalk virus and rotavirus occurred primarily because of the development of immunoassay techniques to recognize these viruses in stool samples and to measure antibodies to them in infected individuals. For Norwalk virus, these assays are currently available in only a few research laboratories (3,4) including that of the principal investigator. This is because the procedure requires use of precious limited human volunteer materials (stools and sera). The assay has more recently been made more efficient in detecting Norwalk virus antigen in stools through the use of an enzyme-linked immunoassay (EIA) instead of a radioimmunoassay (RIA) (5). Together with our collaborators, our use of these immunoassays has shown a major role for this virus in producing in the U.S. clam and oyster associated gastroenteritis, as well as some cases of

travelers' diarrhea in Mexico and Thailand (6-8).

As for rotavirus, use of immunoassay techniques to detect the virus is now common and is employed routinely in many clinical diagnostic laboratories (9). More recently, a monoclonal antibody based EIA that we developed for detection of rotavirus (10) has been shown to be more sensitive and specific than polyclonal antibody tests and has eliminated specificity problems with stool samples from young infants. We have used rotavirus immunoassays to establish the role of rotavirus in several nations around the world, including travelers' diarrhea experienced by U.S. military populations overseas (7, 11-15).

The roles of other enteric viruses in gastroenteritis are incompletely understood, and because of the medical importance of infectious diarrhea, there is clearly a major need to establish the significance of different viruses that may be involved. Comparative studies on their occurrence, however, have been infrequent and usually limited to electron microscopy (15-17). The major obstacle in evaluating the relative importance of the non-rotavirus and non-Norwalk virus enteric viruses as causative agents of gastroenteritis has been the lack of convenient methods for their diagnosis. In addition, for appropriate treatment and control measures to be initiated, rapid as well as convenient methods are required, but have been also unavailable for most of these gastroenteritis viruses. Further, many of these viruses are difficult to cultivate or have not been cultivated in cell culture, which has inhibited characterization studies.

Among these agents, the evidence currently seems strongest that enteric adenoviruses and now astroviruses are medically important pathogens like rotavirus and Norwalk virus. The enteric adenoviruses differ from the well characterized conventional serotypes of adenoviruses which are propagated in standard tissue cultures and are not commonly associated with gastroenteritis. These adenoviruses can not be recognized specifically as enteric types by electron microscopy in stools and can be cultivated only inefficiently in an adenovirus transformed cell line, Graham 293 (18). Two enteric serotypes (types 40 and 41) have been identified and in a limited number of studies performed to date, have been associated with gastroenteritis in infants and young children and much less commonly found in asymptomatic children (1,19). The potential role of enteric adenoviruses in travelers' diarrhea or in disease in adults has been little studied. Convenient and specific immunoassays to detect enteric adenoviruses have been only recently developed and now permit an understanding of their epidemiology as has already occurred with the use of immunoassays to study rotavirus and Norwalk virus. Four years ago we prepared monoclonal antibodies specific for adenovirus types 40 and 41. These antibodies were characterized and used in an EIA format to detect the enteric adenoviruses in known positive diarrheal stool specimens with 95 to 98 percent sensitivity and specificity

(20,21). In 1988 we reported that two percent of acute diarrheal episodes among Thai children were due to enteric adenoviruses using the EIA procedure (22). We are using our enteric adenovirus monoclonal antibody EIA to assess the epidemiology of this infection in other populations on a continuing basis.

Astroviruses are small (27-35mm in diameter) round viruses and have been identified by electron microscopy in the stools of some patients with gastroenteritis (1,23). Astroviruses also have been shown to be cultivatable in cell culture (24,25). However, simple diagnostic procedures have not been widely available prior to our work. Thus, the extent of the role of astroviruses in human diarrheal disease is not known. In 1988, we published our confirmation of in vitro cultivation of human astroviruses (26). This permitted us to purify sufficient viral antigen to prepare monoclonal antibodies reactive against a common antigen shared by multiple astrovirus serotypes (26), which offered the practical possibility for developing immunoassays to assess the medical importance of astroviruses in human viral gastroenteritis. Such an EIA test for the detection of astroviruses in human stools was developed by us and was published in the Journal of Infectious Diseases (27). Since that time we have been using the assay extensively in ongoing epidemiological studies.

Caliciviruses have also been associated with diarrheal disease in humans (1,28). These agents are currently detected mostly by electron microscopy and more convenient assays for their detection are needed so that their epidemiology can be studied. Norwalk virus possesses a single structural protein, characteristic of a calicivirus (29), and the two agents are of similar size and general shape (albeit, differing somewhat in virion surface structure). Thus, the possibility of relatedness between these two enteric viruses exists and was studied by us two years ago. We demonstrated that antigenic characteristics are shared between calicivirus and Norwalk virus based on our detection of seroconversions to Norwalk virus in patients experiencing gastroenteritis due to a strain of calicivirus (30, 31). These two agents, therefore, may belong to the same family of viruses, as also may Snow Mountain agent for which we have also found seroconversions to Norwalk virus in some affected patients (32). These serological cross-reactions demonstrate the need for convenient viral antigen specific detection methods for calicivirus such as we previously developed for Norwalk virus.

ASTROVIRUS ETIOLOGY OF VIRAL GASTROENTERITIS

Prior to our studies, astroviruses had been associated with gastroenteritis but had not been directly shown to be etiological agents in a controlled study. In our annual report for 1990 we described two studies in Thailand which indicated a causative role for astroviruses. To further assess the role and medical impor-

tance of astroviruses in gastroenteritis in these studies, we compared the clinical symptoms found in gastroenteritis patients infected only with astroviruses to those infected only with rotaviruses; patients with other bacterial, viral or parasitic infections were excluded. The results are given in Table 1. The frequency of clinical findings detected with each of these two virus infections was similar, and no significant differences were found in any of the categories evaluated. Dehydration ≥ 5 percent was more common with rotavirus infections but was not significant ($P < 0.08$).

The rate of hospitalization was low in both rotavirus and astrovirus infections, but there was a slightly higher percentage of hospitalized children who had rotavirus infection alone, 3.4 percent (6 of 175) compared to 2.2 percent (1 of 44) for those with astrovirus infection alone. This may have been due to the higher rate of rotavirus infected patients showing dehydration ≥ 5 percent (27 of 175 compared to 2 of 44), although all of the remaining patients with astrovirus infections showed a lower level of dehydration (<5 percent) as did the remainder of those infected with rotaviruses. The results of these studies are now in press (New England Journal of Medicine, reference 33).

Table 1. Clinical findings associated with astrovirus and rotavirus gastroenteritis

Clinical finding	% showing clinical finding	
	Astrovirus infection ^a n = 44	Rotavirus infection ^a n = 175
Watery stools	61	67
Loose stools	41	35
Mucoid stools	55	51
Bloody stools	7	6
Nausea	71	88
Abdominal pain	58	63
Vomiting	61	67
Fever	80	83
Dehydration $\geq 5\%$	5	15

^aIncludes only those samples where no bacterial, parasitic, or other viral pathogens were detected.

In addition to correlating clinical findings with astrovirus infection in this population, we are also testing for sets of viruses, bacteria, and parasites that discriminate between patients with and without gastroenteritis by means of the multivariate method, stepwise logistic regression. A summary of the major findings to date are shown in Table 2. The greater the chi-square

improvement value, the greater the independent association of a given pathogen with disease status. It can be seen that astroviruses were ranked quite high compared to some of the more widely recognized pathogens.

Table 2. Ranking of pathogens as discriminators of disease by logistic regression analysis.

Rank	Pathogen	Improvement Chi-square
1	Rotavirus	272.6
2	<u>Shigella</u> spp.	223.4
3	Enteropathogenic <u>E. Coli</u>	40.1
4	Astrovirus	31.5
5	<u>Cryptosporidium</u> spp.	15.7
6	<u>Salmonella</u> spp.	13.0
7	<u>Campylobacter jejuni</u>	10.5

During this contract year, additional studies on astroviruses have been done in a variety of populations. The data given in Table 3 summarizes the major findings to date for astrovirus gastroenteritis in pediatric populations. All show a major medical role for astroviruses.

Table 3. Incidence of astrovirus diarrhea in various populations and comparison to rotavirus and enteric adenovirus infection.

Geographic Area	No. Tested	Diarrheic (+ or -)	% virus positive		
			Rotavirus	Astrovirus	Enteric Adenovirus
Thailand	1,691	+	19.0	8.2	2.6
(2 studies)	1,459	-	1.0	2.0	0.5
Atlanta,	524	+	nt	4.0	nt
Georgia ^a	105	-	nt	1.0	nt
Australia	1,343	+	6.8	5.5	1.3
(Aborigines)					
Guatemala	5,234	+	3.0	3.7	0.2

^a In collaboration with R.I. Glass, CDC, Atlanta, submitted for publication.

CALICIVIRUS MONOCLONAL ANTIBODIES

To maximize the possibility of obtaining monoclonal antibodies with the limited amount of virus we have available, we initially examined methods for purifying virus from stools using feline calicivirus seeded stools as a model, and applied the results of these model studies to purification of calicivirus antigen from human stool samples. These samples were known to contain calicivirus by IEM. Fractions were inoculated into mice for production of antisera and eventual use in hybridomas. By ELISA techniques we determined that virus-specific antibody was being produced by the mice, and fusions for hybridoma production were done. Clones producing antibody to calicivirus, as determined by ELISA, have been obtained and we are in the process of evaluating them. The utility of these monoclonal antibodies in diagnostic tests remains to be determined.

We are also trying to prepare antibodies to calicivirus antigen in rabbits. This would provide us with sera for use in immunoassays along with our monoclonal antibodies. The antigen used as inoculum was purified from stool material, and because we have a limited quantity of material to work with, a recently introduced adjuvant (Titermax, Cytrx Corp, Norcross, GA), which is purported to be superior to Freund's adjuvant, was used.

SNOW MOUNTAIN VIRUS MONOCLONAL ANTIBODIES

Sometime ago, we had received human peripheral blood lymphocytes from volunteer patients who had received Snow Mountain Virus (courtesy of Dr. R. Dolin, University of Rochester). We had transformed these cells with Epstein-Barr virus for use in future studies. To look at the possibility of preparing human monoclonal antibodies with these and other cells, the transformed cells were fused with human x mouse heterohybridomas. These will hopefully result in hybridomas which produce human antibody specific for Snow Mountain virus. We are in the process of preparing and screening such hybridomas.

ANTIIDIOTYPIC ANTIBODIES AS SURROGATE ANTIGENS

To perform diagnostic assays designed to detect viral antigen, a positive control antigen is necessary. This can be a problem if virus cannot be cultivated and there is a limited amount of clinical material available for this purpose. It has been shown for several viruses, such as hepatitis B virus, Herpes simplex virus, and Newcastle Disease virus among others, that anti-idiotypic antibodies to viral antibodies, which have been likened to an "internal image" of an antigen, can be used as a substitute for the antigen itself. To demonstrate feasibility of this

approach, we are working with astrovirus as a model for the non-cultivable viruses (calicivirus, Norwalk virus, SRVs). We have now generated polyclonal anti-idiotypic antibody to astrovirus monoclonal antibody, which demonstrates that this approach is practical and sound. We are also preparing monoclonal anti-idiotypic antibodies to astrovirus monoclonal antibody as a model system for other viruses. To do this, BALB/c mice were inoculated with our BALB/c mouse-derived monoclonal antibody to astrovirus (purified IgG, emulsified in Freund's incomplete adjuvant, given i.p at 75 ug IgG/mouse). Five additional inoculations were given at two week intervals and mice were bled and tested for anti-idiotypic antibody. For this test, the serum was substituted for viral antigen in our astrovirus EIA. Pre-immune serum was used as a control. The positive reactions obtained indicate that monoclonal antibodies can be derived from these reactive mice, and we are now starting to evaluate these materials in monoclonal antibody based assays for detection of astrovirus. If we can successfully obtain a monoclonal antibody based immunoassay for astrovirus diagnosis without the need for cultivated virus, this would bode well for our ability to generate similar assays for calicivirus and for Norwalk virus.

TYPE SPECIFIC ASTROVIRUS IDENTIFICATION

During the contract year, work was initiated to produce type-specific monoclonal antibodies to each of the five known astrovirus serotypes for type-specific identification of these viruses in immunological and epidemiological studies.

Because we have samples obtained in successive years from the same areas which we have shown contain astroviruses at a relatively high incidence 8% of total diarrheal cases, these samples can be tested with serotype specific astrovirus monoclonal antibodies once they are developed. This will give us information on both prevailing serotypes and possible immunity to them. We have started monoclonal antibody production to astrovirus type 1, which has been found to be the most prevalent type in the limited studies which have been done.

COLLABORATIVE EPIDEMIOLOGICAL STUDIES WITH THE MILITARY

We have continued to perform collaborative studies with U.S. military scientists during the current contract year on the role of viral agents in gastroenteritis.

A series of collaborative efforts with Dr. Peter Echeverria (AFRIMS, Thailand) are the outpatient studies of endemic longitudinally studied diarrhea among children in Bangkok. The adenovirus component has been published (22), and the astrovirus studies are now in press (33). Some of the results are discussed in the section above entitled "Astrovirus etiology of viral gastroenteritis".

Astroviruses have caused outbreaks of gastroenteritis in homes for the elderly, but have not been looked for in younger adult populations. Therefore, we examined for astroviruses in samples obtained from U.S. military personnel with gastroenteritis, in this case from those who were on maneuvers in Honduras, and found that 19.4% (7/36) of swabs were virus positive. Use of convalescent serum in an IEM test on one of the swabs tested showed small, round virus particles which was further evidence that the samples were astrovirus positive. This finding suggests that astroviruses can be important in younger adults as well as in children and the elderly.

LITERATURE CITED

1. Cukor, G. and Blacklow, N.R.: Human viral gastroenteritis. Microbiological Reviews 48:157-179, 1984.
2. Echeverria, P., Blacklow, N.R., Cukor, G., Vibulbandhitkit, S., Changchawalit, S., and Boonthai, P.: Rotavirus as a cause of severe gastroenteritis in adults. J.Clin.Microbiol. 18:663-667, 1983.
3. Greenberg, H.B., Wyatt, R.G., Valdesuso, J., Kalica, A.R., London, W.T., Chanock, R.M., and Kapikian, A.Z.: Solid-phase microtiter radioimmunoassay for detection of the Norwalk strain of acute nonbacterial, epidemic gastroenteritis virus and its antibodies J.Med.Virol. 2:97-108, 1978.
4. Blacklow, N.R., Cukor, G., Bedigian, M.K., Echeverria, P., Greenberg, H.B., Schreiber, D.S., and Trier, J.S.: Immune response and prevalence of antibody to Norwalk enteritis virus as determined by radioimmunoassay. J. Clin. Microbiol. 10:903-909, 1979.
5. Herrmann, J.E., Nowak, N.A., and Blacklow, N.R.: Detection of Norwalk virus in stools by enzyme-immunoassay. Journal of Medical Virology 17: 127-133, 1985.
6. Morse, D.L., Guzewich, J.J., Hanrahan, J.P., Stricof, R., Shayegani, M., Deibel, R., Grabau, J.C., Nowak, N.A., Herrmann, J.E., Cukor, C., and Blacklow, N.R.: Widespread outbreaks of clam and oyster associated gastroenteritis: Role of Norwalk virus. New England Journal of Medicine 314:678-681, 1986.
7. Keswick, B.H., Blacklow, N.R., Cukor, G., DuPont, H.L., and Vollet, J.J.: Norwalk virus and rotavirus in travelers' diarrhoea in Mexico. Lancet 1:109-110, 1982.
8. Echeverria, P., Blacklow, N.R., Sanford, L.B., and Cukor, G.: Travelers' diarrhea among American Peace Corps volunteers in rural Thailand. J.Infect.Dis. 143:767-771, 1981.
9. Blacklow, N.R. and Cukor, G.: Viral gastroenteritis agents. Manual of Clinical Microbiology, Fourth Edition, Ed., Lennette, E. et al (American Society for Microbiology) 805-812, 1985.
10. Herrmann, J.E., Blacklow, N.R., Perron, D.M. et al. Enzyme immunoassay with monoclonal antibodies for the detection of rotavirus in stool specimens. J.Infect.Dis. 152:831-32, 1985.
11. Echeverria, P., Ho, M.T., Blacklow, N.R., Quinnan, G., Portnoy, B., Olson, J.G., Conklin, R., DuPont, H. and Cross, J.H.: Relative importance of viruses and bacteria in the etiology of pediatric diarrhea in Taiwan. J. Infect. Dis. 136:383, 1977.
12. Echeverria, P., Blacklow, N.R., Vollet, J.J.: Reovirus-like agent and enterotoxigenic Escherichia coli infections in pediatric diarrhea in the Philippines, J. Infect. Dis. 138:326, 1978.

13. Echeverria, P., Hodge, F.A., Blacklow, N.R., Vollet, J.J., Cukor, G., DuPont, H.L., and Cross, J.H.: Travelers' diarrhea among United States Marines in South Korea. Am. J. Epidemiol. 108:68, 1978.
14. Echeverria, P., Ramirez, G., Blacklow, N.R., Ksiazek, T., Cukor, G., and Cross, J.H.: Travelers' diarrhea among United States Army troops in South Korea. J.Infect.Dis. 139:215, 1979.
15. Echeverria, P., Blacklow, N.R., Zipkin, C., Vollet, J.J., Olson, J.A., DuPont, H.L. and Cross, J.H.: Etiology of gastroenteritis among Americans living in the Philippines. Am. J. Epidemiol. 109:493-501, 1979.
16. Middleton, P.J. Role of viruses in pediatric gastrointestinal disease and epidemiologic factors. In Virus infections of the gastrointestinal tract, Tyrrell DAJ and Kapikian AZ, eds., Marcel Dekker, Inc., New York, p. 221-225, 1982.
17. Riepenhoff-Talty, M., Saif, L.J., Barrett, H.J., Suzuki, H., Ogra, P.L. Potential spectrum of etiological agents of viral enteritis in hospitalized infants. J. Clin. Microbiol. 17:352-6, 1983.
18. Takiff, H.E., Straus, S.E., and Garon, C.F.: Propagation and in vitro studies of previously noncultivable enteric adenoviruses in 293 cells. Lancet 2:832-834, 1981.
19. Uhnoo, I., Wadell, G., Svensson, L., and Johansson, M.: Importance of enteric adenovirus 40 and 41 in acute gastroenteritis in infants and young children. J.Clin.Micro. 20:365-372, 1984.
20. Herrmann, J.E., Perron-Henry, D.M., Stobbs-Walro, D., Blacklow, N.R.: Preparation and characterization of monoclonal antibodies to enteric adenovirus types 40 and 41. Arch. Virol. 94:259-65, 1987.
21. Herrmann, J.E., Perron-Henry, D.M., Blacklow, N.R. Antigen detection with monoclonal antibodies for the diagnosis of adenovirus gastroenteritis. J. Infect. Dis. 155:1167-71, 1987.
22. Herrmann, J.E., Blacklow, N.R., Perron-Henry, D.M., Clements, E., Taylor, D.N. and Echeverria, P.: Incidence of enteric adenoviruses among children in Thailand and the significance of these viruses in gastroenteritis. J.Clin.Micro. 26:1783-1786, 1988.
23. Kurtz, J.B., Lee, T.W., and Pickering, D.: Astrovirus-associated gastroenteritis in a children's ward. J.Clin.Path. 30:948-952, 1977.
24. Lee, T.W., and Kurtz, J.B.: Serial propagation of astrovirus in tissue culture with the aid of trypsin. J.Gen.Virol. 57:421-424, 1981.
25. Lee, T.W., and Kurtz, J.B.: Human astrovirus serotypes. J. Hyg. 89: 539-540, 1982.
26. Herrmann, J.E., Hudson, R.W., Perron-Henry, D.M., Kurtz, J.B., Blacklow, N.R. Antigenic characterization of cell-cultivated astrovirus serotypes and development of astrovirus-specific monoclonal antibodies.

J.Infect.Dis., 158:182-185,1988.

27. Herrmann, J.E., Nowak, N.A., Perron-Henry D.M., Hudson, R.W., Cubitt, W.D. and Blacklow, N.R. 1990. Diagnosis of astrovirus gastroenteritis by antigen detection with monoclonal antibodies. Journal of Infectious Diseases 161:226-229.
28. Cubitt, W.D., McSwiggan, D.A., and Moore, W.: Winter vomiting disease caused by calicivirus. J.Clin.Pathol. 32:786-793, 1979.
29. Greenberg, H.B., Valdesuso, J.R., Kalica, A.R., Wyatt, R.G., McAuliffe, V.J., Kapikian, A.Z. and Chanock, R.M.: Proteins of Norwalk virus. J. Virol. 37:994-999, 1981.
30. Cubitt, W.D., Blacklow, N.R., Herrmann, J.E. et al. Antigenic relationships between human caliciviruses and Norwalk virus. J. Infect. Dis. 156:806-14, 1987.
31. Blacklow, N.R., Herrmann, J.E. and Cubitt, W.D.: Immunobiology of Norwalk virus. In Novel Diarrhoea Viruses, CIBA Foundation Symposium, Number 128, John Wiley Publishers, London, pp. 144-161, 1987.
32. Guest, C., Spitalny, K.C., Madore, H.P., Pray, K., Dolin, R., Herrmann, J.E., and Blacklow, N.R.: Foodborne Snow Mountain agent gastroenteritis in a school cafeteria. Pediatrics 79:559-563, 1987.
33. Herrmann, J.E., Taylor, D.N., Echeverria, P.E., and Blacklow, N.R. 1991, Astroviruses as a cause of Gastroenteritis in children. New Engl. J. Med. 324: 1757-1760.

PUBLICATIONS OF WORK SUPPORTED BY THIS CONTRACT

1. Herrmann, J.E., Perron-Henry, D.M., Stobbs-Walro, D., and Blacklow, N.R.: Preparation and characterization of monoclonal antibodies to enteric adenovirus types 40 and 41. Archives of Virology, 94:259-265, 1987.
2. Herrmann, J.E., Perron-Henry, D.M., and Blacklow, N.R.: Antigen detection with monoclonal antibodies for the diagnosis of adenovirus gastroenteritis. J.Inf.Dis. 155:1167-1171, 1987.
3. Guest, C., Spitalny, K.C., Madore, H.P., Pray, K., Dolin, R., Herrmann, J.E., and Blacklow, N.R.: Foodborne Snow Mountain agent gastroenteritis in a school cafeteria. Pediatrics 79:559-563, 1987.
4. Blacklow, N.R., Herrmann, J.E. and Cubitt, W.D.: Immunobiology of Norwalk virus. In Novel Diarrhoea Viruses, CIBA Foundation Symposium, Number 128, John Wiley Publishers, London, pp. 144-161, 1987.
5. Cubitt, W.D., Blacklow, N.R., Herrmann, J.E., Nowak, N.A., Nakata, S., and Chiba, S.: Antigenic relationships among human caliciviruses and Norwalk virus. Journal of Infectious Diseases 156:806-814, 1987.
6. Hudson, R.W., Perron-Henry, D.M., Herrmann, J.E., and Blacklow, N.R.: Antigenic relationships of cell-cultivated astrovirus serotypes and development of virus-specific monoclonal antibodies. Abstracts of the 27th Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC), p. 99, 1987.
7. Herrmann, J.E., Hudson, R.W., Blacklow, N.R. and Cubitt, W.D.: Marin County agent, an astrovirus. The Lancet 2:743, 1987.
8. Perron-Henry, D., Herrmann, J.E., Blacklow, N.R., Clements, E., Taylor, D.N. and Echeverria, P.: Prevalence of enteric adenoviruses among children in Thailand. Abstracts of the Annual Meeting of the American Society for Microbiology, p. 351, 1988.
9. Hudson, R.W., Herrmann, J.E., Cubitt, W.D. and Blacklow, N.R.: Propagation of the Marin county strain of astrovirus and its detection by monoclonal antibody. Abstracts of the Annual Meeting of the American Society for Microbiology, p. 351, 1988.
10. Herrmann, J.E., Hudson, R.W., Perron-Henry, D.M., Kurtz, J.B., and Blacklow, N.R.: Antigenic characterization of cell-cultivated astrovirus serotypes and development of astrovirus-specific monoclonal antibodies. Journal of Infectious Diseases, 158:182-185, 1988.
11. Perron-Henry, D., Herrmann, J.E., and Blacklow, N.R.: Isolation and propagation of enteric adenoviruses in HEP-2 cells. Abstracts of the 28th Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC), p.197, 1988.
12. Taylor, D.N., Echeverria, P., Pitarangsi, C., Seriwatana, J., Sethabutr,

- O., Bodhidatta, L., Brown, C., Herrmann, J.E., and Blacklow, N.R.: Application of DNA hybridization techniques in the assessment of diarrheal disease among refugees in Thailand. American Journal of Epidemiology 127:179-187, 1988.
13. Echeverria, P., Taylor, D.N., Leksomboon, U., and Blacklow, N.R.: Case-control study of endemic diarrheal disease in Thai children under 5 years old. Abstracts of the 28th Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC), p. 116, 1988.
 14. Perron-Henry, D.M., Herrmann, J.E., and Blacklow, N.R.: Isolation and propagation of enteric adenoviruses in HEp-2 cells. Journal of Clinical Microbiology 26:1445-1447, 1988.
 15. Taylor, D.N., Echeverria, P., Sethabutr, O., Pitarangsi, C., Lexsomboon, U., Blacklow, N.R., Rowe, B., Gross, R., and Cross, J.: Clinical and microbiologic features of Shigella and enteroinvasive Escherichia coli infections detected by DNA hybridization. Journal of Clinical Microbiology 26:1362-1366, 1988.
 16. Herrmann, J.E., Blacklow, N.R., Perron-Henry, D.M., Clements, E., Taylor, D.N. and Echeverria, P.: Incidence of enteric adenovirus among children in Thailand and the significance of these viruses in gastroenteritis. Journal of Clinical Microbiology 26:1783-1786, 1988.
 17. Herrmann, J.E., and Blacklow, N.R.: Gastroenteritis viruses. In Diagnostic Procedures for Viral, Rickettsial and Chlamydial Infections, Sixth Edition, R. Emmons and N. Schmidt, Eds., American Public Health Association, Washington, D.C., 1989.
 18. Fleissner, M.L., Herrmann, J.E., Booth, J.W., Blacklow, N.R. and Nowak, N.A.: Role of Norwalk virus in two foodborne outbreaks of gastroenteritis: Definitive virus association. American Journal of Epidemiology, 129:165-172, 1989.
 19. Echeverria, P., Taylor, D.N., Lexsomboon, U., Bhaibulaya, M., Blacklow, N.R., Tamura, K., and Sakazaki, R.: Case-control study of endemic diarrheal disease in Thai children. Journal of Infectious Diseases 159: 543-558, 1989.
 20. Hudson, R.W., Herrmann, J.E., and Blacklow, N.R.: Plaque quantitation and virus neutralization assays for human astroviruses. Abstracts of the Annual Meeting of the American Society for Microbiology, p. 388, 1989.
 21. Blacklow, N.R., and Herrmann, J.E.: Norwalk virus. In Viruses and the Gut, Proceedings of the Ninth BSG-SK&F International Workshop 1988, M.J.G. Farthing, Ed., Swan Press Ltd, London, pp. 65-69, 1989.
 22. Herrmann, J.E., Nowak, N., Perron-Henry, D., Hudson, R.W., Cubitt, W.D., and Blacklow, N.R.: Antigen detection with monoclonal antibodies for the diagnosis of astrovirus gastroenteritis. Abstracts of the 29th Interscience Conference on Antimicrobial Agents and Chemotherapy, (ICAAC), p. 176, 1989.

23. Hudson, R.W., J.E. Herrmann, and N.R. Blacklow. 1989. Plaque quantitation and virus neutralization assays for human astroviruses. 1989. Abstracts of the American Society for Microbiology, p. 388.
24. Hudson, R.W., J.E. Herrmann, and N.R. Blacklow. 1989. Plaque quantitation and virus neutralization assays for human astroviruses. Archives of Virology 108:33-38.
25. Perron-Henry, D., Herrmann, J.E., Blacklow, N.R., Albert, M.J., and Erlich, J.: Incidence of rotaviruses and enteric adenoviruses in Aboriginal children from central Australia. Abstracts of the Annual Meeting of the American Society for Microbiology, p. 365, 1990.
26. Herrmann, J.E., N.A. Nowak, D.M. Perron-Henry, R.W. Hudson, W.D. Cubitt, and N.R. Blacklow. 1990. Diagnosis of astrovirus gastroenteritis by antigen detection with monoclonal antibodies. Journal of Infectious Diseases 161:226-229.
27. Herrmann, J.E., W.D. Cubitt, R.W. Hudson, D.M. Perron-Henry, L.S. Oshiro and N.R. Blacklow. 1990. Immunological characterization of the Marin County strain of astrovirus. Archives of Virology 110:213-220.
28. Herrmann, J.E., N.A. Nowak, N.R. Blacklow, D. Perron-Henry, R.W. Hudson, E. Clements, D.N. Taylor, and P. Echeverria. 1990. Astrovirus etiology of pediatric gastroenteritis. Abstracts of the 30th Interscience Conference of Antimicrobial Agents and Chemotherapy, Abstract 1221.
29. Herrmann, J.E., and N.R. Blacklow. 1990. Rotavirus. In Principles and Practice of Infectious Diseases, G. Mandell, R.G. Douglas, and J.E. Bennett, Eds., Churchill Livingstone, Inc., New York, Third Edition, pp. 1234-1240.
30. Moe, C.L., J.R. Allen, S.S. Monroe, J.E. Herrmann, N.R. Blacklow, B. Forrester, M. Koch, and R.I. Glass. 1991. Detection of astrovirus by immunoassay and RNA probe in pediatric stool samples. Abstracts of the American Society for Microbiology, abstract C-3.
31. Lew, J.F., C.L. Moe, J.R. Allen, S.S. Monroe, J.C. Hierholzer, J.E. Herrmann, N.R. Blacklow, A.V. Bartlett, and R.I. Glass. 1991. Astrovirus and adenovirus associated with diarrhea in children attending three different day care settings. Abstracts of the Annual Meeting of the American Society for Microbiology, abstract C-4.
32. Matsui, S.M., J.P. Kim, H.B. Greenberg, J.E. Herrmann, L.S. Oshiro, N.R. Blacklow, and G.R. Reyes. 1991. Identification and characterization of a human astrovirus-specific cDNA. Abstracts of the American Gastroenterology Association, p. A597.
33. Monroe, S.S., S.E. Stine, L. Gorelkin, J.E. Herrmann, N.R. Blacklow, and R.I. Glass. 1991. Temporal synthesis of proteins and RNAs during human astrovirus infection of cultured cells. Journal of Virology 65:641-648.

34. Herrmann, J.E., Taylor, D.N., Echeverria, P.E., and Blacklow, N.R.
1991, Astroviruses as a cause of Gastroenteritis in children. N. Engl.
J. Med. 324: 1757-1760.